

CONTENTS

| SL.No | PARTICULARS | PAGE NO |
|--------------|----------------------------------------------------|----------------|
| 1. | Preface | 2 |
| 2. | Methodology | 3 |
| 3. | Methodology Protocol for Marine water analysis | 4 |
| 4. | Methodology Protocol for Sediment quality analysis | 5 |
| 5. | Location map of sampling stations | 8 |
| 6. | Sampling Stations With GPS | 9 |
| 7. | Physical Properties-Water | 11 |
| 8. | Chemical Properties-Water | 12 |
| 9. | Nutrients - Water | 13 |
| 10. | Heavy Metals - Water | 14 |
| 11. | Biological Characteristics | 15 |
| 12. | Phytoplankton | 16-17 |
| 13. | Zooplankton | 18-19 |
| 14. | Sediments Quality Data - Nutrients | 21 |
| 15. | Texture - Sediment | 22 |
| 16. | Sediments Quality Data - Heavy Metals | 23 |
| 17. | Macrobenthos- Sediment | 24-25 |
| 18. | Meiobenthos-Sediment | 26-27 |
| 19. | Standards | 28 |

PREFACE

Environmental monitoring programme is a vital process of any management plan of a development project. Concern over the state of environment has grown worldwide since the sixties, due to decline in environmental quality, and various efforts have been taken for environmental protection in our country. Accordingly, the Ministry of Environment & Forests, Govt. of India, became the nodal agency in regulating developmental activities enforcing environmental sampling and monitoring.

Dredging Corporation of India Ltd (DCI), is one among the Public Sector Undertakings of India, provides dredging services to the Major Ports of the country in India and is a pioneer organization in the field of dredging and maritime development. Mormugao Port Trust (MPT), Goa, entrusted the work of Capital Dredging of the approach channel, turning circle, berths 5,6,7 and approach for capsized vessels at Mormugao port, Goa.

Dredging Corporation of India Ltd (DCI), Visakhapatnam took the services from **M/s. Richardson & Cruddas (1972) Ltd, Chennai-98(A Govt. of India Undertaking)**, for environmental monitoring in and around the dredging and dumping areas of Mormugao port through their **Work order No. DCI/HSE/IMS/28 dtd. 19.02.2016**. Accordingly, the sample of marine water and sediment during dredging was collected on: **07.04.2016**. The samples collected during dredging were analysed and presented in this report. **The analysis data reveals that the marine water and sediment quality is well within the standards prescribed by Ministry of Environment and Forest (MoEF).**

Grateful thanks are due to **Dr. P.K.Sethi, Joint General Manager (HSE)** and all other supporting staff of **Dredging Corporation of India Ltd (DCI)** for the opportunity provided to be associated in this endeavor.

Place: Chennai

Date: 22.04.2016

(E.BALAKRISHNAIAH)

Unit In-charge

METHODOLOGY

SAMPLING METHODOLOGY:

Marine Water

Marine Water samples were collected using a bottom sampler. On-site test such as pH, salinity, Temp., EC, Turbidity etc. were carried out immediately after the sample collection. The samples intended for chemical, heavy metal and bacteriological analyses are preserved with necessary reagents and analysed in the laboratory. The plankton samples were collected using plankton net of diameter of 0.35 m, No.25 mesh size 63 μ . The plankton net was towed for 15 minute at the sampling locations for collection of samples for estimation of Phytoplankton and Zooplankton.

The Parameter covered are:-

Physical Properties: pH, EC, Colour, Odour, Salinity, Temperature, Turbidity, TSS

Chemical Properties: DO, COD, BOD, Oil & Grease, Nutrients, Sulphates, Chlorides

Heavy Metals : Fe, Zn, Mg, Cd, Cr, Hg

Marine Biology : primary productivity, Chlorophyll and Phytoplankton & Zooplankton

Sediment

Marine sediment samples were collected using a Peterson's Grab Sampler. The collected sediment samples were segregated on the site for analysis of physico-chemical parameters, heavy metals and benthic communities. The sediment sample for benthic communities subject to sieving for recording the macro benthos and then the samples and preserved with Rose Bengal and Formalin Solution for further analysis of Benthic communities

The Parameter covered are:

Physico-chemical Properties: Texture, pH, Organic Matter, Nutrients, Oil and Grease.

Heavy Metals : Fe, Mn, Cd, Ni, Cr, Hg, Zn and Pb

Benthic Communities : Macro & Micro Benthic Flora and Fauna

METHODOLOGY PROTOCOL FOR MARINE WATER ANALYSIS

| S.No. | Parameters | Methodology Protocol |
|------------------------------|------------------------------------------|-------------------------------|
| Physical properties | | |
| 1 | pH | IS 3025 Part 11 (Reaff. 2006) |
| 2 | Colour | IS 3025 Part 4 (Reaff. 2006) |
| 3 | Odour | IS 3025 Part 5 (Reaff. 2006) |
| 4 | Electrical Conductivity | IS 3025 Part 14 (Reaff. 2006) |
| 5 | Temperature | IS 3025 Part 9 (Reaff. 2006) |
| 6 | Salinity | APHA 22nd Edn. 2520 |
| 7 | Turbidity | IS 3025 Part 10 (Reaff. 2006) |
| 8 | Total Suspended Solids | IS 3025 Part 17 (Reaff. 2006) |
| Chemical properties | | |
| 9 | Dissolved Oxygen | IS 3025 Part 38 (Reaff. 2009) |
| 10 | COD | APHA 22st Edn. 5220 B |
| 11 | BOD-3 Days, 27°C | APHA 22st Edn. 5210 B |
| 12 | Oil & Grease | IS 3025 Part 39 (Reaff. 2009) |
| 13 | Chlorides (as Cl) | IS 3025 Part 32 (Reaff. 2009) |
| 14 | Fluorides (as F) | IS 3025 Part 60 (Reaff. 2008) |
| 15 | Sulphates (as SO ₄) | IS 3025 Part 24 (Reaff. 2009) |
| 16 | Total Nitrogen (as N) | IS 3025 Part 34 (Reaff. 2009) |
| 17 | Nitrate Nitrogen (as NO ₃ -N) | IS 3025 Part 34 (Reaff. 2009) |
| 18 | Total Phosphate (as PO ₄ -P) | IS 3025 Part 31 (Reaff. 2009) |
| Heavy metals | | |
| 19 | Iron | APHA 22nd Edn. 3500-Fe |
| 20 | Zinc | APHA 22nd Edn. 3500-Zn |
| 21 | Magnesium | APHA 22nd Edn. 3500-Mg |
| 22 | Cadmium | APHA 22nd Edn. 3500-Cd |
| 23 | Chromium | APHA 22nd Edn. 3500-Cr |
| 24 | Mercury | APHA 22nd Edn. 3500-Hg |
| Biological parameters | | |
| 25 | Phyto & Zoo Planktons and Pigments | APHA 22nd Edn. 10200 |

METHODOLOGY PROTOCOL FOR SEDIMENT QUALITY ANALYSIS

| S.No. | Parameters | Methodology Protocol |
|----------------------------|-----------------|-------------------------------|
| Physical properties | | |
| 1 | pH | IS 3025 Part 11 (Reaff. 2006) |
| 2 | Organic matters | IS 2720 Part 22 (Reaff. 1995) |
| 3 | Nutrients | IS 10158 -1982 |
| 4 | Oil & Grease | IS 3025 Part 39 (Reaff. 2009) |
| Heavy metals | | |
| 5 | Iron | EPA 7380 |
| 6 | Manganese | EPA 7460 |
| 7 | Cadmium | EPA 7130 |
| 8 | Nickel | EPA 7520 |
| 9 | Chromium | EPA 7090 |
| 10 | Mercury | EPA 7471 B |
| 11 | Zinc | EPA 7950 |
| 12 | Lead | EPA 7420 |
| Benthic Communities | | |
| 13 | Macro benthos | APHA 22nd Edn. 10500 |
| 14 | Meio benthos | APHA 22nd Edn. 10700 |

Monitoring and Testing of Marine water & Sediment samples for Capital Dredging inside the Mormugao Port, Goa.

Summary Report

Marine water and sediment samples were collected in seven stations at Mormugao Port, as per the locations identified by the DCI. The survey made on **07.04.2016** for dredging phase.

Physico-chemical parameters such as Temperature, Colour, Odour, Salinity, pH, Dissolved oxygen, COD, BOD, Turbidity, Total Suspended Solids, Chlorides, Sulphates, nutrients and Heavy metals were estimated by standard methods. Biological variables have also been studied and this includes Phytoplankton, Zooplankton and its Biomass. Sediment samples were collected and analyzed the pH, Total Organic Carbon, Total Phosphorus, Total Nitrogen, Soil Texture, Heavy metals and Macro and Meio benthos.

The observations made during this period revealed the following information which has been grouped in terms of three variables such as physical, chemical and biological. The sea surface temperature varied between 28.0°C to 30.0°C and there was no significant variation in temperature with the distance from the shore. The salinity ranged from 30.06 to 33.21‰. The pH of the seawater samples observed from 8.08 to 8.56. The measured turbidity varied between 10 to 18 NTU. The TSS value varied from 12 to 28mg. The concentration of cadmium in water was found to be <0.001mg/l. The chromium values was found to be <0.001mg/l, Ferrous from 0.50 to 0.75 mg/l, Magnesium from 1524 to 1724 mg/l and Zinc from 0.32 to 0.46 mg/l. The concentration of mercury shows the BDL (<0.001mg/l) level. The population density of Phytoplankton varied from 4198 to 6280 Cell/L. The higher phytoplankton density was recorded at station **SPOIL GROUND -I**, The species such as, *Coscinodiscus eccentricus*, *Coscinodiscus centralis*, *Chaetoceros affinis*, *Pleurosigma normanii*, *Cerataulina orientalis* and *Thalassionema nitzschiaoides* were

found to be common in all stations monitored. The numerical abundance of zooplankton varied from 3870 to 5760 Organisms/m³. The higher zooplankton density was recorded at station **SPOIL GROUND -I**. Zooplankton consists of Paracalanus parvus, Oithona similis, Corycaeas danae, Favella philipiensis, Copepod nauplii and Sagitta sp were found to be dominant species commonly distributed in all the stations monitored.

The concentrations of Ferrous in sediments were ranging from 4068 to 5742 µg/g. Manganese from 32.62 to 48.62 µg/g. Cadmium in sediments ranged between 0.48 to 0.86 µg/g. Nickel from 1.78 to 2.68 µg/g. The chromium varied from 12.42 to 17.93 µg/g. The concentration of mercury varied from 0.15 to 0.39 µg/g. The concentrations of Zinc varied from 19.99 to 27.68 µg/g and the Lead from 11.56 to 15.92 µg/g. The numerical abundance of the macro benthic fauna varied from 1520 to 2530 No/square meter and the Meiofauna varied between 175 to 255 No/10cm²

Concluding Remarks

As per the Env. Monitoring made during **dredging phase (07.04.2016)** suggests the following conclusion

- The marine water quality at 7 locations were found to be well within the primary water quality criteria for class SW - IV waters (Harbour water)
- The sediment quality at 7 locations were found to be well within the hazardous waste management rules 2003 (schedule 2)

**POSITIONS OF PRE DETERMINED LOCATIONS FOR SEA WATER / SEDIMENT SAMPLE
AT MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No: | Nomenclature | in UTM | | in Geo-graphic | |
|----------------------|----------------------------|------------------|-----------------|-----------------------|-----------------|
| | | NORTHINGS | EASTINGS | Lat (N) | Long (E) |
| DUMPING AREA | | | | | |
| 1 | SPOIL GROUND -II | 1707641 | 356430 | 15° 26' 30".56 | 73° 39' 42".75 |
| 2 | SPOIL GROUND -I | 1707690 | 358512 | 15° 26' 32".57 | 73°40' 52".57 |
| DREDGING AREA | | | | | |
| 3 | A0 | 1703771 | 363378 | 15° 24' 26".01 | 73° 43' 36".60 |
| 4 | A1 | 1704176 | 365421 | 15° 24' 39".59 | 73° 44' 45".05 |
| 5 | A2 | 1704391 | 366966 | 15° 24' 46".87 | 73° 45' 36".84 |
| 6 | A3 | 1704780 | 368843 | 15° 24' 59".87 | 73° 46' 39".72 |
| 7 | Between A4 & A5 | 1705091 | 370377 | 15° 25' 10.27 | 73° 47' 31".12 |

Marine Water Quality data

PHYSICAL PROPERTIES

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No. | Sample description | pH | Colour (Hazen unit) | Odour | EC (micro mhos/cm) | W.T (°C) | Salinity (ppt) | Turbidity (NTU) | TSS (mg/l) |
|----------------------|----------------------------|------|---------------------------|-----------|-----------------------|-------------|-------------------|--------------------|---------------|
| DUMPING AREA | | | | | | | | | |
| 1 | SPOIL GROUND -II | 8.56 | 9 | Odourless | 52300 | 28.0 | 33.21 | 10 | 12 |
| 2 | SPOIL GROUND -I | 8.24 | 12 | Odourless | 51200 | 28.5 | 31.26 | 16 | 20 |
| DREDGING AREA | | | | | | | | | |
| 3 | A0 | 8.10 | 8 | Odourless | 50100 | 29.5 | 31.52 | 10 | 14 |
| 4 | A1 | 8.20 | 8 | Odourless | 52600 | 29.5 | 32.10 | 10 | 14 |
| 5 | A2 | 8.08 | 14 | Odourless | 50100 | 29.0 | 30.06 | 18 | 28 |
| 6 | A3 | 8.18 | 10 | Odourless | 53900 | 29.5 | 31.14 | 12 | 16 |
| 7 | Between A4 & A5 | 8.11 | 16 | Odourless | 51200 | 30.0 | 32.10 | 16 | 24 |

CHEMICAL PROPERTIES –WATER

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No. | Sample description | DO (mg/l) | COD (mg/l) | BOD (mg/l) | Oil & Grease (mg/l) | Chloride (mg/l) | Sulphate (mg/l) |
|---------|----------------------------|--------------|---------------|---------------|---------------------------|--------------------|--------------------|
| 1. | SPOIL GROUND -II | 5.2 | 79 | 1 | <1 | 19682 | 3192 |
| 2. | SPOIL GROUND -I | 5.0 | 72 | 2 | <1 | 18962 | 2975 |
| 3. | A0 | 4.6 | 73 | 2 | <1 | 18762 | 3274 |
| 4. | A1 | 5.1 | 84 | 2 | <1 | 19786 | 3412 |
| 5. | A2 | 4.8 | 75 | 1 | <1 | 18735 | 3284 |
| 6. | A3 | 5.3 | 78 | 2 | <1 | 19984 | 3451 |
| 7. | Between A4 & A5 | 4.9 | 75 | 1 | <1 | 18956 | 2969 |

NUTRIENTS – WATER

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| S. No. | Station Code | Parameters (mg/l) | | | |
|--------|----------------------------|-------------------|----------------|-----------------|------------------|
| | | Amm.Nitrogen | Total Nitrogen | Total Phosphate | SiO ₂ |
| 1. | SPOIL GROUND -II | 1.5 | 3.5 | 1.4 | 20.9 |
| 2. | SPOIL GROUND -I | 1.1 | 2.3 | 1.2 | 19.0 |
| 3. | A0 | 1.2 | 2.4 | 1.1 | 19.2 |
| 4. | A1 | 1.6 | 3.4 | 1.3 | 19.5 |
| 5. | A2 | 1.2 | 2.5 | 1.2 | 18.9 |
| 6. | A3 | 1.7 | 3.8 | 1.5 | 21.5 |
| 7. | Between A4 & A5 | 1.2 | 2.9 | 1.2 | 19.6 |

HEAVY METALS – WATER

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No. | Station Code | Parameter (mg/l) | | | | | |
|---------|----------------------------|------------------|------|------|--------|--------|--------|
| | | Fe | Zn | Mg | Cd | Cr | Hg |
| 1. | SPOIL GROUND -II | 0.62 | 0.42 | 1697 | <0.001 | <0.001 | <0.001 |
| 2. | SPOIL GROUND -I | 0.50 | 0.32 | 1524 | <0.001 | <0.001 | <0.001 |
| 3. | A0 | 0.54 | 0.33 | 1552 | <0.001 | <0.001 | <0.001 |
| 4. | A1 | 0.65 | 0.39 | 1652 | <0.001 | <0.001 | <0.001 |
| 5. | A2 | 0.59 | 0.37 | 1604 | <0.001 | <0.001 | <0.001 |
| 6. | A3 | 0.75 | 0.46 | 1724 | <0.001 | <0.001 | <0.001 |
| 7. | Between A4 & A5 | 0.55 | 0.35 | 1562 | <0.001 | <0.001 | <0.001 |

BIOLOGICAL CHARACTERISTICS

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| S. No. | Station Code | Chl a (mg/m ³) | Phaeopigment (mg/m ³) | Net Primary Productivity (mg C/ m ³ /d) |
|--------|----------------------------|-------------------------------|--------------------------------------|----------------------------------------------------------|
| 1 | SPOIL GROUND -II | 2.83 | 0.78 | 0.25 |
| 2 | SPOIL GROUND -I | 2.23 | 0.64 | 0.23 |
| 3 | A0 | 2.58 | 0.62 | 0.20 |
| 4 | A1 | 2.87 | 0.81 | 0.27 |
| 5 | A2 | 2.65 | 0.72 | 0.22 |
| 6 | A3 | 3.06 | 0.89 | 0.28 |
| 7 | Between A4 & A5 | 2.55 | 0.60 | 0.21 |

PHYTOPLANKTON

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No | Species (Cells/l) | Location ID | | | |
|--------|-------------------------------------|------------------|-----------------|-------------|-------------|
| | | SPOIL GROUND -II | SPOIL GROUND -I | A0 | A1 |
| | Bacillariophyceae | | | | |
| 1. | <i>Bacteriastrum comosum</i> | 160 | 250 | 280 | 110 |
| 2. | <i>Cerataulina orientalis</i> | 145 | 220 | 150 | 160 |
| 3. | <i>Chaetoceros affinis</i> | 160 | 230 | 320 | 140 |
| 4. | <i>Chaetoceros indicus</i> | 110 | 220 | 240 | 200 |
| 5. | <i>Coscinodiscus centralis</i> | 160 | 210 | 240 | 180 |
| 6. | <i>Coscinodiscus eccentricus</i> | 210 | 240 | 260 | 250 |
| 7. | <i>Coscinodiscus granii</i> | 240 | 250 | 180 | 180 |
| 8. | <i>Coscinodiscus gigas</i> | 120 | 220 | 240 | * |
| 9. | <i>Ditylum brightwelli</i> | 230 | 370 | 240 | 210 |
| 10. | <i>Gyrosigma balticum</i> | * | 280 | 310 | 220 |
| 11. | <i>Leptocylindrus danicus</i> | * | 240 | 250 | 205 |
| 12. | <i>Lithodesmium undulatum</i> | 260 | 220 | 240 | 250 |
| 13. | <i>Odontella mobiliensis</i> | 310 | 360 | 110 | 110 |
| 14. | <i>Pleurosigma normanii</i> | 260 | 210 | 260 | 110 |
| 15. | <i>Skeletonema costatum</i> | 150 | 200 | 240 | 210 |
| 16. | <i>Stephanophysis palmeriana</i> | 300 | 290 | 310 | * |
| 17. | <i>Thalassionema nitzschiooides</i> | 230 | 310 | 110 | 220 |
| 18. | <i>Thalassiothrix frauenfeldii</i> | * | 210 | 210 | 120 |
| 19. | <i>Triceratium favus</i> | 150 | 200 | 200 | * |
| 20. | <i>Triceratium reticulatum</i> | 180 | 160 | 310 | 220 |
| | Cyanophyceae | | | | |
| 21. | <i>Anabeana nastoc</i> | 90 | 110 | 120 | 210 |
| 22. | <i>Microcystis sp.</i> | 70 | 240 | 280 | 110 |
| 23. | <i>Trichodesmium erythraeum</i> | 170 | 210 | 320 | 150 |
| 24. | <i>Rhizosolenia styliformis</i> | 240 | 290 | * | * |
| | Dinoflagellates | | | | |
| 25. | <i>Ceratium furca</i> | 180 | 180 | 250 | 280 |
| 26. | <i>Ceratium macroceros</i> | 260 | 150 | 240 | 180 |
| 27. | <i>Ceratium tripos</i> | 220 | 210 | 250 | 250 |
| | Total | 4605 | 6280 | 6160 | 4275 |

* - Organisms not present

PHYTOPLANKTON

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No | Species (Cells/l) | A2 | A3 | Between A4 & A5 |
|--------|-------------------------------------|-------------|-------------|-----------------|
| | Bacillariophyceae | | | |
| 1. | <i>Bacteriastrum comosum</i> | 110 | 145 | 210 |
| 2. | <i>Cerataulina orientalis</i> | 220 | 224 | 320 |
| 3. | <i>Chaetoceros affinis</i> | 230 | 208 | 210 |
| 4. | <i>Chaetoceros indicus</i> | 220 | 180 | 180 |
| 5. | <i>Coscinodiscus centralis</i> | 210 | 110 | 250 |
| 6. | <i>Coscinodiscus eccentricus</i> | 240 | 200 | 200 |
| 7. | <i>Coscinodiscus granii</i> | 110 | 110 | 310 |
| 8. | <i>Coscinodiscus gigas</i> | 220 | 205 | 205 |
| 9. | <i>Ditylum brightwelli</i> | 220 | 150 | * |
| 10. | <i>Gyrosigma balticum</i> | 280 | 110 | 250 |
| 11. | <i>Leptocylindrus danicus</i> | 240 | 150 | 350 |
| 12. | <i>Lithodesmium undulatum</i> | 220 | 110 | 260 |
| 13. | <i>Odontella mobiliensis</i> | 250 | 120 | 120 |
| 14. | <i>Pleurosigma normanii</i> | 210 | 100 | 340 |
| 15. | <i>Skeletonema costatum</i> | 200 | 246 | 320 |
| 16. | <i>Stephanophysis palmeriana</i> | 240 | 195 | 310 |
| 17. | <i>Thalassionema nitzschiooides</i> | 310 | 250 | 250 |
| 18. | <i>Thalassiothrix frauenfeldii</i> | 210 | 110 | 110 |
| 19. | <i>Triceratium favus</i> | 200 | * | 250 |
| 20. | <i>Triceratium reticulatum</i> | 160 | 90 | 310 |
| | Cyanophyceae | | | |
| 21. | <i>Anabeana nastoc</i> | 110 | 100 | 310 |
| 22. | <i>Microcystis sp.</i> | 240 | 150 | 150 |
| 23. | <i>Trichodesmium erythraeum</i> | 110 | 140 | 140 |
| 24. | <i>Rhizosolenia alata</i> | 150 | 90 | 90 |
| 25. | <i>Rhizosolenia styliformis</i> | 180 | 110 | 110 |
| | Dinoflagellates | | | |
| 26. | <i>Ceratium furca</i> | 150 | 210 | 210 |
| 27. | <i>Ceratium macroceros</i> | 210 | 110 | 110 |
| 28. | <i>Ceratium tripos</i> | 200 | 110 | 310 |
| 29. | <i>Protoperdinium oceanicum</i> | 250 | 165 | * |
| | Total | 5900 | 4198 | 6185 |

* - Organisms not present

ZOOPLANKTONSample Collected at: **MORMUGAO PORT, GOA**Sample Collected on: **07.04.2016**

| Sl. No | Species (Organisms/m ³) | Location ID | | | |
|--------------|-------------------------------------|------------------|-----------------|-------------|-------------|
| | | SPOIL GROUND -II | SPOIL GROUND -I | A0 | A1 |
| | Copepoda | | | | |
| 1 | <i>Acartia spinicauda</i> | 210 | 250 | * | * |
| 2 | <i>Acartia erythrea</i> | * | 210 | 210 | 190 |
| 3 | <i>Acrocalanus gipper</i> | * | 220 | 190 | * |
| 4 | <i>Acrocalanus gracilis</i> | 190 | 150 | * | 220 |
| 5 | <i>Centropages furcatus</i> | 250 | 250 | 270 | * |
| 6 | <i>Nannocalanus minor</i> | 190 | 150 | 300 | 250 |
| 7 | <i>Paracalanus parvus</i> | 210 | 190 | 280 | 140 |
| 8 | <i>Pontella danae</i> | 150 | 160 | * | 200 |
| 9 | <i>Temora turbinata</i> | 200 | 220 | 210 | 180 |
| 10 | <i>Oithona brevicornis</i> | 250 | 320 | 180 | 140 |
| 11 | <i>Oithona rigida</i> | 180 | 210 | * | * |
| 12 | <i>Oithona similis</i> | 200 | 210 | 210 | 180 |
| 13 | <i>Corycaeas danae</i> | 110 | 180 | 270 | 220 |
| 14 | <i>Copilia mirabilis</i> | 100 | 220 | 210 | 180 |
| | Spirotricha | | | | |
| 15 | <i>Favella brevis</i> | 200 | 180 | 250 | 200 |
| 16 | <i>Favella philipiensis</i> | 180 | 220 | 220 | 180 |
| 17 | <i>Tintinnopsis tubulosa</i> | * | 190 | 290 | 110 |
| 18 | <i>Tintinnopsis tocantinensis</i> | 110 | 320 | * | * |
| 19 | <i>Tintinnopsis cylinderica</i> | 180 | 160 | 190 | 100 |
| | Others | | | | |
| 20 | <i>Lucifer hansperi</i> | 210 | 240 | * | 200 |
| 21 | <i>Sagitta sp</i> | 180 | 220 | 160 | 320 |
| 22 | <i>Oikopleura dioica</i> | * | 300 | * | 140 |
| 23 | <i>Oikopleura parva</i> | 210 | 180 | 210 | 160 |
| | Larval Forms | | | | |
| 24 | <i>Bivalve Veliger</i> | 200 | 210 | 110 | 210 |
| 25 | <i>Barnacle nauplii</i> | 160 | 220 | 210 | * |
| 26 | <i>Copepod nauplii</i> | 110 | 180 | 200 | 220 |
| 27 | <i>Crustacean nauplii</i> | 140 | 200 | 250 | 200 |
| Total | | 4120 | 5760 | 4420 | 3940 |

* - Organisms not present

ZOOPLANKTON

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No | Species (Organisms/m ³) | Location ID | | |
|--------|-------------------------------------|-------------|-------------|-----------------|
| | | A2 | A3 | Between A4 & A5 |
| | Copepoda | | | |
| 1 | <i>Acartia spinicauda</i> | * | 110 | 260 |
| 2 | <i>Acartia erythrea</i> | 250 | 250 | 240 |
| 3 | <i>Acrocalanus gipper</i> | 210 | * | * |
| 4 | <i>Acrocalanus gracilis</i> | 180 | 220 | * |
| 5 | <i>Centropages furcatus</i> | 220 | 110 | 250 |
| 6 | <i>Nannocalanus minor</i> | * | 180 | * |
| 7 | <i>Paracalanus parvus</i> | 260 | 210 | 210 |
| 8 | <i>Pontella danae</i> | * | 180 | 200 |
| 9 | <i>Temora turbinata</i> | 210 | * | 260 |
| 10 | <i>Oithona brevicornis</i> | * | 140 | 240 |
| 11 | <i>Oithona rigida</i> | 180 | * | 190 |
| 12 | <i>Oithona similis</i> | 250 | 210 | * |
| 13 | <i>Corycaeas danae</i> | 210 | 150 | 290 |
| 14 | <i>Copilia mirabilis</i> | 280 | * | 250 |
| | Spirotricha | | | |
| 15 | <i>Favella brevis</i> | * | 180 | * |
| 16 | <i>Favella philipiensis</i> | 330 | 210 | 290 |
| 17 | <i>Tintinnopsis tubulosa</i> | 230 | 210 | 190 |
| 18 | <i>Tintinnopsis tocantinensis</i> | * | 180 | * |
| 19 | <i>Tintinnopsis cylinderica</i> | 180 | 210 | 180 |
| | Others | | | |
| 20 | <i>Lucifer hansperi</i> | 260 | 200 | * |
| 21 | <i>Sagitta sp</i> | 190 | 150 | 190 |
| 22 | <i>Oikopleura dioica</i> | * | * | 280 |
| 23 | <i>Oikopleura parva</i> | 230 | 240 | 360 |
| | Larval Forms | | | |
| 24 | <i>Bivalve Veliger</i> | 210 | 180 | 190 |
| 25 | <i>Barnacle nauplii</i> | 190 | * | * |
| 26 | <i>Copepod nauplii</i> | 220 | 210 | 220 |
| 27 | <i>Crustacean nauplii</i> | * | 140 | 180 |
| | Total | 4290 | 3870 | 4470 |

* - Organisms not present

SEDIMENT

Quality data

pH, NUTRIENTS & TOTAL ORGANIC CARBON, OIL & GREASE – SEDIMENT

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| S. No. | Station Code | pH | Total Nitrogen ($\mu\text{g/g}$) | Total Phosphorus ($\mu\text{g/g}$) | Total Organic Carbon (mg/g) | O & G ($\mu\text{g/g}$) |
|--------|----------------------------|-----|------------------------------------|--------------------------------------|-----------------------------|---------------------------|
| 1. | SPOIL GROUND -II | 8.4 | 13.46 | 9.05 | 4.25 | 0.682 |
| 2. | SPOIL GROUND -I | 8.5 | 11.45 | 8.25 | 3.59 | 0.526 |
| 3. | A0 | 8.3 | 12.49 | 8.56 | 3.85 | 0.529 |
| 4. | A1 | 8.5 | 13.58 | 9.12 | 4.70 | 0.723 |
| 5. | A2 | 8.3 | 12.59 | 8.86 | 4.08 | 0.576 |
| 6. | A3 | 8.6 | 13.96 | 9.25 | 4.95 | 0.762 |
| 7. | Between A4 & A5 | 8.5 | 12.13 | 8.45 | 3.86 | 0.595 |

TEXTURE – SEDIMENT

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| S. No. | Station Code | Grain Size Distribution (%) | | |
|--------|----------------------------|-----------------------------|------|------|
| | | Sand | Silt | Clay |
| 1. | SPOIL GROUND -II | 3.2 | 19.4 | 77.4 |
| 2. | SPOIL GROUND -I | 3.1 | 18.5 | 78.4 |
| 3. | A0 | 7.2 | 20.5 | 72.3 |
| 4. | A1 | 8.5 | 19.5 | 72.0 |
| 5. | A2 | 7.5 | 20.9 | 71.6 |
| 6. | A3 | 8.5 | 21.6 | 69.9 |
| 7. | Between A4 & A5 | 9.1 | 21.2 | 69.7 |

HEAVY METALS – SEDIMENT

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No. | Station Code | $\mu\text{g/g}$ | | | | | | | |
|------------|----------------------------|-----------------|-------|------|------|-------|------|-------|-------|
| | | Fe | Mn | Cd | Ni | Cr | Hg | Zn | Pb |
| 1. | SPOIL GROUND –II | 4068 | 35.56 | 0.59 | 1.78 | 12.75 | 0.24 | 19.99 | 12.95 |
| 2. | SPOIL GROUND -I | 4258 | 43.45 | 0.86 | 2.68 | 17.93 | 0.19 | 27.68 | 15.92 |
| 3. | A0 | 4562 | 34.26 | 0.48 | 2.06 | 14.28 | 0.26 | 22.56 | 12.95 |
| 4. | A1 | 5692 | 45.62 | 0.64 | 2.45 | 14.06 | 0.28 | 25.62 | 14.62 |
| 5. | A2 | 4958 | 38.25 | 0.49 | 2.18 | 12.69 | 0.22 | 23.56 | 14.26 |
| 6. | A3 | 5742 | 48.62 | 0.65 | 2.64 | 14.23 | 0.39 | 27.50 | 15.62 |
| 7. | Between A4 & A5 | 4235 | 32.62 | 0.65 | 1.99 | 12.42 | 0.15 | 20.62 | 11.56 |

MACROBENTHOS DISTRIBUTION IN THE SEDIMENT

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No | Species (No/m ²) | Location ID | | | |
|--------------|------------------------------|------------------|-----------------|-------------|-------------|
| | | SPOIL GROUND -II | SPOIL GROUND -I | A0 | A1 |
| | Polychaetes | | | | |
| 1 | <i>Armandia longicaudata</i> | 90 | 110 | 120 | 150 |
| 2 | <i>Capitella capitata</i> | 110 | 100 | * | 130 |
| 3 | <i>Cirriformia sp</i> | 110 | 120 | 110 | 150 |
| 4 | <i>Goniada emerita</i> | 110 | * | 150 | * |
| 5 | <i>Nephtys dibranchis</i> | * | 150 | 210 | 130 |
| 6 | <i>Nereis sp.</i> | 90 | * | 120 | * |
| 7 | <i>Notomastus aberans</i> | * | 200 | 100 | 110 |
| 8 | <i>Perinereis capensis</i> | 100 | * | 100 | 180 |
| 9 | <i>Platynereis calodonta</i> | 110 | * | 210 | 210 |
| 10 | <i>Prionospio cirrifera</i> | 200 | * | * | * |
| 11 | <i>Prionospio pinnata</i> | * | 140 | 100 | 110 |
| | Bivalves | | | | |
| 12 | <i>Donax veligers</i> | 120 | * | * | * |
| 13 | <i>Meretrix veligers</i> | * | 120 | * | 240 |
| | Gastropods | | | | |
| 14 | <i>Littorina veligers</i> | 100 | * | 110 | 160 |
| 15 | <i>Natica veligers</i> | 120 | 100 | * | 120 |
| 16 | <i>Nassarius variegatus</i> | 100 | 50 | 110 | 110 |
| 17 | <i>Turris veligers</i> | 110 | 110 | 210 | 150 |
| | Crustaceans | | | | |
| 18 | <i>Ampithoe romondi</i> | 120 | 90 | 100 | * |
| 19 | <i>Angeliera phreaticola</i> | 120 | 110 | 150 | 250 |
| 20 | <i>Gynodiastylis sp.</i> | 110 | * | * | 110 |
| 21 | <i>Paragnathia formica</i> | 110 | 120 | 110 | 110 |
| Total | | 1930 | 1520 | 2010 | 2420 |

* - Organisms not present

MACROBENTHOS DISTRIBUTION IN THE SEDIMENT

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No | Species (No/m ²) | Location ID | | |
|--------------------|------------------------------|-------------|-------------|-------------|
| | | A2 | A3 | Between |
| Polychaetes | | | | |
| 1 | <i>Armandia longicaudata</i> | 140 | 110 | 90 |
| 2 | <i>Capitella capitata</i> | 110 | 150 | 120 |
| 3 | <i>Cirriformia sp</i> | * | * | 110 |
| 4 | <i>Goniada emerita</i> | 100 | 130 | 210 |
| 5 | <i>Nephtys dibranchis</i> | 150 | 120 | 90 |
| 6 | <i>Nereis sp.</i> | 110 | 110 | * |
| 7 | <i>Notomastus aberans</i> | 220 | 130 | * |
| 8 | <i>Perinereis capensis</i> | 110 | 220 | 90 |
| 9 | <i>Platynereis calodonta</i> | 110 | 110 | 110 |
| 10 | <i>Prionospio cirrifera</i> | 90 | 100 | 100 |
| 11 | <i>Prionospio pinnata</i> | 110 | 210 | 80 |
| Bivalves | | | | |
| 12 | <i>Donax veligers</i> | 100 | 90 | 90 |
| 13 | <i>Meretrix veligers</i> | * | 100 | 80 |
| Gastropods | | | | |
| 14 | <i>Littorina veligers</i> | * | 210 | 110 |
| 15 | <i>Natica veligers</i> | 110 | 110 | 90 |
| 16 | <i>Nassarius variegatus</i> | 220 | * | * |
| 17 | <i>Turris veligers</i> | 180 | * | * |
| Crustaceans | | | | |
| 18 | <i>Ampithoe romondi</i> | 110 | 180 | 90 |
| 19 | <i>Angeliera phreaticola</i> | 90 | 210 | 110 |
| 20 | <i>Gynodiastylis sp.</i> | 80 | 110 | 110 |
| 21 | <i>Paragnathia formica</i> | 100 | 130 | 120 |
| Total | | 2240 | 2530 | 1800 |

* - Organisms not found

MEIOBENTHOS distribution in the sediment

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No | Species (No/10cm ²) | Location ID | | | |
|--------------|---------------------------------|------------------|------------------|------------|------------|
| | | SPOIL GROUND -II | SPOIL GROUND - I | A0 | A1 |
| | Foraminiferans | | | | |
| 1 | <i>Ammonia beccarii</i> | 15 | 10 | 5 | 10 |
| 2 | <i>Bolivina sp.</i> | 19 | 5 | 15 | 15 |
| 3 | <i>Cibicides refulgens</i> | 12 | 7 | * | * |
| 4 | <i>Globorotalia hiruste</i> | 13 | * | 6 | 15 |
| 5 | <i>Loxostomum sp.</i> | 10 | * | 9 | 12 |
| 6 | <i>Miliammina sp.</i> | 18 | 8 | 21 | 17 |
| 7 | <i>Milionella sp.</i> | * | 15 | 20 | 14 |
| 8 | <i>Nonion sp</i> | 10 | 7 | 18 | 14 |
| | Nematodes | | | | |
| 9 | <i>Daptonema conicum</i> | * | * | 11 | 16 |
| 10 | <i>Draconema sp.</i> | 10 | 14 | 11 | 10 |
| 11 | <i>Greeffiella sp.</i> | * | 11 | 7 | 12 |
| 12 | <i>Microlaimus sp.</i> | 18 | 12 | 18 | 14 |
| 13 | <i>Neochromodora sp.</i> | 15 | 10 | 14 | 12 |
| 14 | <i>Spirinia sp.</i> | 10 | * | * | 5 |
| 15 | <i>Synonchus sp.</i> | * | * | 10 | 10 |
| 16 | <i>Theristus sp.</i> | 10 | 16 | 20 | 10 |
| 17 | <i>Viscosia sp.</i> | 14 | 7 | 9 | 18 |
| | Ostrocodes | | | | |
| 18 | <i>Cypridies sp.</i> | 9 | 10 | * | * |
| 19 | <i>Cytheromorpha sp.</i> | * | 9 | 10 | 8 |
| 20 | <i>Neocytheideis sp.</i> | 15 | 15 | 9 | * |
| 21 | <i>Tanella indica</i> | * | 19 | 9 | 16 |
| 22 | <i>Tanella kingmai</i> | 10 | * | * | 10 |
| Total | | 208 | 175 | 222 | 238 |

* - Organisms not present

MEIOBENTHOS distribution in the sediment

Sample Collected at: **MORMUGAO PORT, GOA**

Sample Collected on: **07.04.2016**

| Sl. No | Species (No/10cm ²) | Location ID | | |
|-----------------------|---------------------------------|-------------|------------|--------------|
| | | A2 | A3 | Between A4 & |
| Foraminiferans | | | | |
| 1 | <i>Ammonia beccarii</i> | 10 | 10 | 9 |
| 2 | <i>Bolivina sp.</i> | 15 | 9 | 5 |
| 3 | <i>Cibicides refulgens</i> | * | 14 | 6 |
| 4 | <i>Globorotalia hiruste</i> | 18 | * | 15 |
| 5 | <i>Loxostomum sp.</i> | * | * | 7 |
| 6 | <i>Miliammina sp.</i> | 20 | 9 | 8 |
| 7 | <i>Milionella sp.</i> | * | 12 | 14 |
| 8 | <i>Nonion sp</i> | 21 | 15 | 12 |
| Nematodes | | | | |
| 9 | <i>Daptonema conicum</i> | * | 12 | 6 |
| 10 | <i>Draconema sp.</i> | 10 | 21 | 16 |
| 11 | <i>Greeffiella sp.</i> | 15 | 12 | 6 |
| 12 | <i>Microlaimus sp.</i> | 14 | 21 | * |
| 13 | <i>Neochromodora sp.</i> | 17 | 24 | 10 |
| 14 | <i>Spirinia sp.</i> | * | * | 15 |
| 15 | <i>Synonchus sp.</i> | * | * | 11 |
| 16 | <i>Theristus sp.</i> | 18 | 16 | 10 |
| 17 | <i>Viscosia sp.</i> | 12 | 18 | 7 |
| Ostrocodes | | | | |
| 18 | <i>Cypridies sp.</i> | 10 | * | * |
| 19 | <i>Cytheromorpha sp.</i> | * | 14 | 10 |
| 20 | <i>Neocytheideis sp.</i> | 22 | 16 | 14 |
| 21 | <i>Tanella indica</i> | * | 12 | 9 |
| 22 | <i>Tanella kingmaii</i> | 10 | 20 | * |
| Total | | 212 | 255 | 190 |

* - Organisms not present

Standards

- 1. Marine water**
- 2. Hazardous waste Management and Handling Rules 2003 – List of waste and Concentration Limits**

Marine Water Quality Standards

Primary Water Quality Criteria for Class SW-IV Waters (For Harbour Waters)

| S.No. | Parameter | Standards | Rationale/Remarks |
|-------|----------------------------------------------------------------------|----------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------|
| 1. | pH range | 6.5-9.0 | To minimize corrosive and scaling effect. . |
| 2. | Dissolved Oxygen | 3.0 mg/l or 40 percent saturation value, which ever is higher. | Considering bio-degradation of oil and inhibition to is oxygen production through photosynthesis. |
| 3. | Colour and Odour | No noticeable colour or offensive odour. | None from reactive chemicals which may corrode paints/metallic surfaces. |
| 4. | Floating Matters Oil, grease and scum (including Petroleum products) | 10 mg/l | Floating matter should be free from excessive living organisms, which may clog or coat operative parts of marine vessels/equipment. . |
| 5. | Fecal Coliform | 500/100 ml (PAN) | Not exceeding 1000/100 ml in 20 percent of samples in the year and in 3 consecutive samples in monsoon months. |
| 6. | Biochemical Oxygen Demand (3 days at 27°C) | 5 mg/l | To maintain water relatively free from pollution caused by sewage and other decomposable wastes |
| 7. | Biochemical Oxygen Demand (BOD) (3 days at 27°C) | 3 mg/l | Restricted for bathing (aesthetic quality of water). Also prescribed by IS:2296 1974. |

Source : EPA, 1986
 [GSR 7, dated Dec. 22, 1998]

Hazardous waste Management and Handling Rules 2003

SCHEDULE - 2 [See rule 3(i) (b)] LIST OF WASTE SUBSTANCES WITH CONCENTRATION LIMITS

Classes

Class A

Concentration limit: 50 mg/kg

- A1 Antimony and antimony compounds
- A2 Arsenic and arsenic compounds
- A3 beryllium and cadmium compounds
- A4 Cadmium and beryllium compounds
- A5 Chromium (VI) compounds
- A6 Mercury and mercury compounds
- A7 Selenium and selenium compounds
- A8 Tellurium and tellurium compounds
- A9 Thallium and thallium compounds
- A10 Inorganic cyanide compounds (cyanides)
- A11 Metal carbonyls
- A12 Napthalene
- A13 Anthracene
- A14 Phenanthrene
- A15 Chrysene, benzo(a) anthracene, fluoranthene, benzo(a) pyrene, benzo(K)fluoranthene, indeno(1, 2, 3-ed) pyrene and benzo(ghi)perylene
- A16 Halogenated fused aromatic rings, e.g. polychlorobiphenyls plus derivatives
- A17 Halogenated aromatic compounds
- A18 Benzene
- A19 Dieldrin, aldrin, and endrin
- A20 Organotin compounds

Class B

Concentration limit: 5,000 mg/kg

- B1 Chromium (III) compounds
- B2 Cobalt compounds
- B3 Copper compounds
- B4 Lead and lead compounds
- B5 Molybdenum compounds
- B6 Nickel compounds
- B7 Tin compounds
- B8 Vanadium compounds
- B9 Tungsten compounds

- B10 Silver compounds
- B11 Organic halogen compounds
- B12 Organic phosphorus compounds
- B13 Organic peroxides
- B14 Organic nitro-and nitroso-compounds
- B15 Organic azo-and azo-oxy compounds
- B16 Nitriles
- B17 Amines
- B18 (Iso-and thio-) cyanates
- B19 Phenol and phenolic compounds
- B20 Merceptans
- B21 Asbestos
- B22 Drilling, cutting, grinding and rolling oil or emulsions thereof
- B23 Halogen-silanes
- B24 Hydrazine(s)
- B25 Fluorine
- B26 Chlorine
- B27 Bromine
- B28 White phosphorus
- B29 Ferro-silicon and alloys
- B30 Manganese-silicon
- B31 Halogen-containing substances which produce acidic vapours on contact with damp air or water, e.g. silicon tetrachloride, aluminum chloride, titanium tetrachloride

Class C

Concentration limit: 20,000 mg/kg

- C1 Ammonia and ammonium compounds
- C2 Inorganic peroxides
- C3 Barium compounds, except barium sulphate
- C4 Fluorine compounds
- C5 Phosphorus compounds, except the phosphates of aluminum, calcium and iron
- C6 Bromates, (hypo)bromites
- C7 Chlorates, (hypo)chlorites
- C8 Aromatic compounds
- C9 Organic silicon compounds
- C10 Organic sulphur compounds
- C11 Iodates
- C12 Nitrates, nitrites
- C13 Sulphides
- C14 Zinc compounds
- C15 Salts of per-acids
- C16 Acid halides, acid amides
- C17 Acid anhydrides

Class D

Concentration limit: 50,000 mg/kg

- D1 Sulphur
- D2 Inorganic acids
- D3 Metal bisulphates
- D4 Oxides and hydroxides except those of: hydrogen, carbon, silicon, iron, aluminum, titanium, manganese, magnesium, calcium
- D5 Aliphatic and napthenic hydrocarbons
- D6 Organic oxygen compounds
- D7 Organic nitrogen compounds
- D8 Nitrides
- D9 Hydrides

Class E

Regardless of concentration limit

- E.1 Highly flammable substances
- E.2 Substances which generate dangerous quantities of highly flammable gases on contact with water or damp air.

* All on dry weight basis